

# Memory T and memory B cells share a transcriptional program of self-renewal with long-term hematopoietic stem cells

Chance John Luckey<sup>\*†</sup>, Deepta Bhattacharya<sup>\*†‡</sup>, Ananda W. Goldrath<sup>\*†§</sup>, Irving L. Weissman<sup>‡</sup>, Christophe Benoist<sup>\*¶</sup>, and Diane Mathis<sup>\*¶</sup>

<sup>\*</sup>Joslin Diabetes Center; Departments of Pathology and Medicine, Brigham and Women's Hospital, Harvard Medical School, 1 Joslin Place, Boston, MA 02215; and <sup>†</sup>Stanford Institute for Stem Cell Biology and Regenerative Medicine, Department of Pathology, Stanford University School of Medicine, Stanford, CA 94305-5323

Contributed by Diane Mathis, December 23, 2005

The only cells of the hematopoietic system that undergo self-renewal for the lifetime of the organism are long-term hematopoietic stem cells and memory T and B cells. To determine whether there is a shared transcriptional program among these self-renewing populations, we first compared the gene-expression profiles of naïve, effector and memory CD8<sup>+</sup> T cells with those of long-term hematopoietic stem cells, short-term hematopoietic stem cells, and lineage-committed progenitors. Transcripts augmented in memory CD8<sup>+</sup> T cells relative to naïve and effector T cells were selectively enriched in long-term hematopoietic stem cells and were progressively lost in their short-term and lineage-committed counterparts. Furthermore, transcripts selectively decreased in memory CD8<sup>+</sup> T cells were selectively down-regulated in long-term hematopoietic stem cells and progressively increased with differentiation. To confirm that this pattern was a general property of immunologic memory, we turned to independently generated gene expression profiles of memory, naïve, germinal center, and plasma B cells. Once again, memory-enriched and -depleted transcripts were also appropriately augmented and diminished in long-term hematopoietic stem cells, and their expression correlated with progressive loss of self-renewal function. Thus, there appears to be a common signature of both up- and down-regulated transcripts shared between memory T cells, memory B cells, and long-term hematopoietic stem cells. This signature was not consistently enriched in neural or embryonic stem cell populations and, therefore, appears to be restricted to the hematopoietic system. These observations provide evidence that the shared phenotype of self-renewal in the hematopoietic system is linked at the molecular level.

Self-renewal is a process by which a daughter cell that maintains the same properties as its parent is generated. The best-studied self-renewing cells are long-term hematopoietic stem cells (Lt-HSC), which maintain themselves as a population for the lifetime of the organism. However, self-renewal within the hematopoietic system is not limited to stem cells, because antigen-specific memory B and T cells have also been observed to self-renew in perpetuity. Although this phenotypic similarity has been noted previously (1–3), there is to date no information on whether these cells use the same molecular pathways for self-renewal. Although the extracellular signals involved in cellular homeostasis likely differ between memory and stem cells, we hypothesized that these external cues converge on some of the common cell-intrinsic mediators involved in self-renewal, perhaps through the reactivation of genetic programs used by Lt-HSC.

Adult Lt-HSC are multipotent cells capable of both lifelong self-renewal and differentiation into the various mature cellular components of blood (4). Differentiation of Lt-HSC leads to the formation of short-term hematopoietic stem cells (St-HSC). Although St-HSC retain full hematopoietic differentiation potential, they have a more limited, “short-term,” self-renewal potential. St-HSC subsequently differentiate into lineage-committed precursors (LCP) of either the myeloid or lymphoid lineages. Further

differentiation of LCP is restricted to their respective lineage, and they are incapable of self-renewal. The inability to undergo self-renewal holds true for all subsequent downstream precursor populations as well as for the majority of mature blood cells. Thus, the self-renewal of Lt-HSC is required for sustained hematopoiesis over the course of an organism's life.

Memory T and B cells are mature blood cells that reacquire the ability to undergo long-term self-renewal and are the product of a carefully controlled process of differentiation in response to immunostimulation, such as infection by pathogens (1–3, 5, 6). Before infection, antigen-inexperienced, or “naïve,” cells of a particular specificity exist at very low frequencies and rarely, if ever, divide (7–9). Upon antigenic exposure, naïve cells capable of recognizing one of the pathogen's components undergo a process of rapid clonal expansion and differentiation. For T cells, this process leads to the generation of effector cells that have acquired the functional capacity to rapidly combat foreign pathogens. Effector T cells undergo a dramatic contraction in numbers after pathogen clearance, with 90–95% of them succumbing to apoptosis within weeks after the initial infection (2, 5). However, a subset of the antigen-specific cells persists long after antigen exposure and constitutes the memory T cell compartment.

For B cells, the early thymus-dependent responses to antigenic challenge lead to the formation of rapidly proliferating, short-lived, antibody-secreting plasma cells and germinal center B cells, which undergo somatic hypermutation and Ig isotype switching. Similar to effector T cells, the vast majority of these two cell types is eliminated through apoptosis (10, 11). The surviving antigen-specific B cells comprise two separate memory compartments: the long-lived antibody-secreting plasma cell and the self-renewing memory B cell. The antibody-secreting plasma B cells are completely quiescent and secrete antigen-specific Ig indefinitely, irrespective of antigen re-exposure (12). In contrast, self-renewing memory B cells proliferate slowly and rapidly respond to antigen reexposure by differentiating into both plasma and germinal center B cells in another round of affinity maturation (10, 13).

Memory lymphocytes respond more robustly than their naïve counterparts to antigenic challenge. This ability to respond, combined with their increased frequency and self-renewal, ensures that reexposure to a particular pathogen leads to rapid and vigorous

Conflict of interest statement: No conflicts declared.

Abbreviations: LCP, lineage-committed precursors; Lt-HSC, long-term hematopoietic stem cells; St-HSC, short-term hematopoietic stem cells.

Data deposition: All data sets have been deposited in the Gene Expression Omnibus, [www.ncbi.nlm.nih.gov/geo](http://www.ncbi.nlm.nih.gov/geo) (accession nos. GSE 4142 and GSM 94871–94882).

<sup>†</sup>C.J.L., D.B., and A.W.G. contributed equally to this work.

<sup>§</sup>Present address: Division of Biology Section of Molecular Biology, University of California at San Diego, 9500 Gilman Drive, San Diego, CA 92093-0377.

<sup>¶</sup>To whom correspondence should be addressed. E-mail: [cbdm@joslin.harvard.edu](mailto:cbdm@joslin.harvard.edu).

© 2006 by The National Academy of Sciences of the USA



renewal function in all three progenitor populations. In addition, because St-HSC retain full differentiation potential, the enriched transcripts are unlikely to represent genes solely involved in lymphocyte biology or fate commitment.


We also looked at those transcripts selectively down-regulated in memory CD8<sup>+</sup> T cells relative to both naïve and effector cells (Fig. 1 E and F). One hundred two transcripts were identified with expression levels selectively down-regulated in memory CD8<sup>+</sup> cells relative to both naïve and effector cells. Again, we observed a strong correlation between expression trends in memory CD8<sup>+</sup> T cells and Lt-HSC. The transcripts that were absent or down-regulated in memory CD8<sup>+</sup> cells were also relatively depleted in Lt-HSC (Fig. 1G). The transcripts most down-regulated in Lt-HSC versus LCP also appeared to be progressively up-regulated as cells differentiated into St-HSC and LCP (Fig. 1H in red). Together with the data presented in Fig. 1 A–D, this result demonstrated that the majority of T cell memory enriched transcripts are also enriched in Lt-HSC, whereas the majority of T cell memory depleted transcripts are depleted in Lt-HSC.

Next we addressed whether a similar correlation could be observed between Lt-HSC and the other self-renewing mature lymphocyte population, memory B cells. Thus, we turned to an independently generated collection of Affymetrix GeneChip data on memory, naïve, germinal center, and plasma B cells. To generate antigen-specific B cells *in vivo*, we immunized mice with the T-dependent immunogen NP-CGG, and antigen-specific cells were harvested at various time-points (detailed in *Supporting Methods*, which are published as supporting information on the PNAS web site). These cells were functionally verified by transfer into RAG<sup>-/-</sup> hosts and measuring T-dependent antibody production (Fig. 6, which is published as supporting information on the PNAS web site).

In a fashion parallel to the above analysis, we compared the gene-expression profiles of memory and naïve B cells (Fig. 2A). The transcripts whose levels were most increased in memory cells were selected and subsequently plotted versus germinal center cells (Fig. 2B) and then plasma cells (Fig. 2C). Thereby, a set of 272 transcripts enriched in memory B cells relative to both naïve, germinal center, and plasma B cells were delineated. Analysis of these enriched transcripts within the various stem-cell populations revealed a preferential representation in Lt-HSC-enriched genes compared with either LCP or St-HSC, both in numbers (71% and 79%, respectively) and *P* values (Fig. 2D). Just as was observed in the T cell analysis, the degree of skewing of the memory B cell-enriched transcripts correlated inversely with the progressive loss of self-renewal capacity (Fig. 2E in red). Clearly then, a large fraction of those transcripts augmented in memory B cells relative to naïve, germinal center, and plasma cells were also selectively enriched in Lt-HSC.

A set of 481 transcripts selectively down-regulated in memory B cells relative to the other B cell populations was also delineated (Fig. 2 F–H). Again, a strong correlation between expression in memory B cells and Lt-HSC was observed, as transcripts depleted in memory B cells were diminished in Lt-HSC relative to both LCP and St-HSC (Fig. 2I). Those transcripts most down-regulated in Lt-HSC versus LCP also appeared to be progressively up-regulated as cells differentiated into St-HSC and LCP (Fig. 2J in red). These observations demonstrate that the majority of B cell-memory-enriched genes were augmented in Lt-HSC, whereas the majority of B cell-memory-depleted transcripts were depleted in Lt-HSC. This separate B cell data set provides an important independent confirmation of the T cell data comparisons.

Those transcripts whose expression was up-regulated in both memory T and memory B cells relative to their non-self-renewing counterparts were tabulated (Fig. 3A). Virtually all (92% and 85%) of these shared transcripts were enriched in Lt-HSC (Fig. 3B). Similar results were obtained when those genes down-regulated in both memory populations were compared, with 88% and 84% also



**Fig. 2.** Memory B cell-enriched transcripts are also enriched in Lt-HSC, whereas memory B cell-depleted transcripts are also depleted in Lt-HSC. The 272 transcripts whose expression was relatively enriched (fold change  $\geq 1.4$ ) in memory B cells relative to naïve (A), germinal center (B), and plasma B cells (C) are plotted for their relative expression in Lt-HSC vs. LCP, Lt-HSC vs. St-HSC, and St-HSC vs. LCP (D). The number of transcripts whose  $\log_2$  (fold change) is greater or less than zero is shown at the top of the plot. Those transcripts whose expression in Lt-HSC vs. LCP was  $> 1.4$  were then highlighted in red (E), and their number greater or less than zero is shown at the bottom of the plot. The 481 transcripts whose expression was relatively depleted ( $\leq -1.4$ ) in memory B cells relative to naïve (F), germinal center (G), and plasma B cells (H) are plotted for their relative expression in Lt-HSC vs. LCP, Lt-HSC vs. St-HSC, and St-HSC vs. LCP (I). The number of transcripts whose  $\log_2$  (fold change) is greater or less than zero is shown at the top of the plot. Those transcripts whose expression in Lt-HSC vs. LCP was  $< -1.4$  were then highlighted in red (J), and their number greater or less than zero is shown at the bottom of the plot.

down-regulated in Lt-HSC (Fig. 3C). We propose that these transcripts underlie the most restrictive transcriptional definition of immune memory, and virtually all of them were coordinately regulated in Lt-HSC. The transcripts expressed in concert among memory T, memory B, and Lt-HSC likely represent a transcriptional profile of self-renewal in these diverse hematolymphoid cells.

Q-PCR provided confirmation of the expression of several of those transcripts coordinately enriched in memory T cells and Lt-HSC. Lt-HSC were purified as Lin<sup>-lo</sup>, Sca<sup>+</sup>, c-kit<sup>+</sup>, CD34<sup>-</sup>, and Flt3<sup>-</sup> cells (19–21). Likewise, St-HSC were purified as Lin<sup>-lo</sup>, Sca<sup>+</sup>, c-kit<sup>+</sup>, CD34<sup>+</sup>, and Flt3<sup>-</sup> cells and LCP as Lin<sup>-lo</sup>, Sca<sup>+</sup>, c-kit<sup>+</sup>, CD34<sup>+</sup>, and Flt3<sup>+</sup> cells. Although this LCP population differs from the one used in the GeneChip analysis in its Sca expression, both populations lack self-renewal capacity and show a limited differentiation capacity (14, 19–22). In nearly every individual comparison, the Q-PCR data confirmed what was observed in the microarray analysis (Fig. 7, which is published as supporting information on the PNAS web site). Indeed, the differences observed by Q-PCR were often much greater than those estimated from the chip data. Two noticeable exceptions were the cytokine receptors IL-18R and IL-7R. IL-18R did not appear to be enriched in either HSC population, whereas IL-7R transcript levels were high in Lt-HSC, low in St-HSC, and highest in LCP.


To observe how these shared transcripts partitioned in other stem cell comparisons, we considered our T cell data in con-

### A Transcripts Up-regulated in *Tmem* and *Bmem*

Symbol	Name	Unigene	Positive Fold Change		
			Lt-HSC vs. LCP	Lt-HSC vs. St-HSC	St-HSC vs. LCP
Plekha5	Pleckstrin homology domain containing, family A	Mm.80531	2.60	1.88	1.38
B930046C15R	RIKEN cDNA B930046C15 gene	Mm.204	2.30	1.65	1.39
Mapk12	Mitogen-activated protein kinase 12	Mm.38343	2.07	1.55	1.34
BC042396	cDNA sequence BC042396	Mm.24944	2.03	1.39	1.45
Il18r1	Interleukin 18 receptor 1	Mm.4773	1.96	1.29	1.53
Il7r	Interleukin 7 receptor	Mm.389	1.73	1.87	0.93
ApoE	Apolipoprotein E	Mm.138866	1.54	1.19	1.30
Sema4f	Semaphorin 4f	Mm.42035	1.26	1.12	1.13
Pou6f1	POU domain, class 6, transcription factor 1	Mm.28825	1.22	1.07	1.15
Prkcz	Protein kinase C, zeta	Mm.28561	1.10	1.10	1.00
Ebi2	Epstein-Barr virus induced gene 2	Mm.3007	1.09	1.09	1.01
na	Similar to zinc finger protein 187	Mm.24023	1.08	1.23	0.88
Usp48	Ubiquitin specific protease 48	Mm.21824	0.90	1.04	0.86
Fcgr2b	Fc receptor, IgG, low affinity IIb	Mm.10809	0.23	0.73	0.32

### Transcripts Down-regulated in *Tmem* and *Bmem*

Symbol	Name	Unigene	Negative Fold Change		
			Lt-HSC vs. LCP	Lt-HSC vs. St-HSC	St-HSC vs. LCP
2310008M10R	RIKEN cDNA 2310008M10 gene	Mm.9811	3.48	1.16	3.01
A1317230	EST A1317230	Mm.151329	2.97	1.34	2.20
Coq7	Demethyl-Q 7	Mm.20634	2.62	1.55	1.69
Ywhah	Eta polypeptide	Mm.3308	2.57	1.52	1.68
Myb	Myeloblastosis oncogene	Mm.1202	2.39	0.75	3.20
Eif1ay	Eukaryotic translation initiation factor 1A, Y-linked	Mm.65264	2.23	1.40	1.60
1110004P21Ri	RIKEN cDNA 1110004P21 gene	Mm.43213	2.23	0.78	2.87
3110003A17Ri	RIKEN cDNA 3110003A17 gene	Mm.28149	2.13	1.18	1.80
Cbx3	Chromobox homolog 3 (Drosophila HP1 gamma)	Mm.28148	1.83	1.53	1.19
Sfrs9	Splicing factor, arginine/serine rich 9	Mm.29426	1.66	1.23	1.35
Snrpe	Small nuclear ribonucleoprotein E	Mm.27669	1.63	1.21	1.35
Hnrpa2b1	Heterogeneous nuclear ribonucleoprotein A2/B1	Mm.16767	1.43	1.58	0.91
Gnb1	Guanine nucleotide binding protein, beta 1	Mm.2344	1.37	1.03	1.33
Erh	Enhancer of rudimentary homolog (Drosophila)	Mm.21952	1.35	2.11	0.64
Hnrpa3	Heterogeneous nuclear ribonucleoprotein A3	Mm.21968	1.02	0.99	1.03
2310061N23Ri	RIKEN cDNA 2310061N23 gene	Mm.46382	0.87	1.10	0.79
Hmrt12	Heterogeneous nuclear ribonucleoproteins	Mm.27545	0.58	2.85	0.21



**Fig. 3.** Coordinately regulated transcripts in both memory CD8<sup>+</sup> T cells and memory B cells are also coordinately regulated in Lt-HSC. (A) Those transcripts whose expression in coordinately regulated in memory B cells and memory T cells are listed. (B) Enriched transcripts are plotted for their relative expression in Lt-HSC vs. LCP, Lt-HSC vs. St-HSC, and St-HSC vs. LCP. (C) Depleted transcripts are plotted for their relative expression in Lt-HSC vs. LCP, Lt-HSC vs. St-HSC, and St-HSC vs. LCP. The number of transcripts whose log<sub>2</sub> (fold change) is greater or less than zero is shown at the top of the plots.

junction with other array comparisons of stem cell populations. Although a number of additional hematopoietic stem cell global gene expression studies have been performed (23–25), previous work has shown that cross-platform comparisons do not correlate well (26). Therefore, we focused our analyses on studies performed with Affymetrix-based experiments and used the complete data sets published by Ivanova *et al.* (14), Ramalho-Santos *et al.* (27), and Akashi *et al.* (28). We are aware of only one other recently published Affymetrix data set comparing HSC populations (29), but we were unable to obtain the original files in time for inclusion in this publication. Collectively, the three independent data sets analyzed included adult and fetal hematopoietic stem cells as well as embryonic and neural stem cells. The vast majority of those genes whose expression we identified as coenriched in memory T cells and memory B cells (*vis-à-vis* their designated counterparts) were also augmented in the adult hematopoietic stem cell populations of Ramalho-Santos *et al.* (27) and Akashi *et al.* (ref. 28; Fig. 4). Furthermore, most of the shared transcripts were also increased in fetal-liver hematopoietic cell precursors of Ivanova *et al.* (14). These findings provide independent confirmation of our results, suggesting that this common “self-renewal” molecular signature might be a general feature of hematopoietic stem cell populations. However, there was not a consistent enrichment of the shared transcripts in either neural stem cell or ES cell populations, arguing that this particular molecular signature may be restricted to the self-renewing cells of the hematopoietic system.


### Discussion

We sought to provide biological evidence for or against the hypothesis that memory T and memory B cells have reacquired the expression of molecules characteristic of long-term stem cells, coincident with their ability to self-renew. The data presented demonstrate that for both memory T and B cells, a significant subset

of their transcripts was also found in Lt-HSC. Indeed, virtually all of those selected transcripts whose expression was most closely coordinated in B and T memory cells were similarly regulated in Lt-HSC. These observations provide evidence supporting the hypothesis that the self-renewal pathways used in memory T and B cells are related to those of hematopoietic stem cells.

Although nearly all of the transcripts shared between memory B and T cells were also found in Lt-HSC, there were many more transcripts shared between only one memory population and Lt-HSC (Tables 1–3, which are published as supporting information on the PNAS web site). Because hematopoietic stem cells are absolutely critical for the survival of the organism, it is likely that they rely on redundant pathways involved in self-renewal, only some of which are used in a given memory lymphocyte population. One explanation for the limited overlap observed between memory T and memory B cells is that memory T cells have reactivated different self-renewal pathways than memory B cells. Support for this explanation can be found by looking at those transcripts shared between Lt-HSC and only one memory population. *Bmi-1* is conserved between memory B cells and Lt-HSC but is not enriched in memory T cells. *Bmi-1* is a polycomb family member involved in self-renewal of hematopoietic stem cells (30), leukemia cells (31), and central and peripheral nervous system cells (32). Conversely, memory T cells and Lt-HSC, but not memory B cells, have up-regulated *Iex-1* and *Spi-2A*. These transcripts are known regulators of apoptosis that function in memory CD8<sup>+</sup> T cell survival (33–35). These examples support the hypothesis that a given memory cell lineage may have reactivated only a subset of the redundant pathways expressed in HSC.

Still other transcripts shared between one memory population and Lt-HSC have functions consistent with their potentially playing a role in self-renewal. Memory T cells and Lt-HSC share expression of TNF receptor II (p74R) and TNF receptor-associated factor 1 (Traf-1). These proteins associate within the cell, resulting in a



**Fig. 4.** Coordinately regulated transcripts in memory B and memory T cells are coordinately regulated in several different data sets of hematopoietic stem cells but are not coordinately regulated in neural or ES cells. Transcripts listed in Fig. 5 are on the y axis, and their expression is plotted as a heat map in each of the comparisons listed on the x axis. Up-regulated transcripts are shown in red, and down-regulated transcripts are shown in green. LiLt and LiSt represent Lt-HSC and St-HSc in the published data of Akashi *et al.* (28) HSC and BM represent the Lt-HSC and mature bone marrow of Ramalho-Santos *et al.* (27). FIH and FIL represent the fetal liver HSC and lineage committed progenitors of Ivanova *et al.* (14). NSC and LVB represent the neural stem cells and lateral ventricular of the brain of Ramalho-Santos *et al.* (27) iNSC represent the neural stem cells of Ivanova *et al.* (14). ESC and BM represent the ES cells and bone marrow of Ramalho-Santos *et al.* (27) iESC and Mbc represent the ES cells and mature bone marrow of Ivanova *et al.* (14). All of the data sets were pooled together for rma analysis as described in *Methods*.

signal that inhibits apoptosis (36, 37). Also present on the shared memory T cell list were several members of the RAS/mitogen-activated protein kinase pathway, known to be involved in decisions by stem cells to undergo proliferation, apoptosis, and differentiation (38–40). Likewise, the memory B cells and Lt-HSC share expression of several classes of transcripts that likely function in self-renewal. Tcf4 and Tcf12 are potentially downstream of  $\beta$ -catenin signaling, itself known to play a role in self-renewal in several stem cell systems (41). Finally, Mef2a and Mef2d are members of a class of transcription factors known to help translate calcium signals in neurons into long-term survival (42, 43). Taken together, the presence of these particular transcripts supports the general hypothesis that memory cells have selectively reactivated different self-renewal molecular pathways found in Lt-HSC.

Even though our data point to different self-renewal pathways being reactivated in either memory B or T cells, we observed several transcripts whose expression was shared between Lt-HSC and both memory populations. Of these jointly shared transcripts, only IL-7R has been shown to play a role in memory T cell self-renewal (5). Although IL-7R clearly plays a role in B cell progenitor differentiation, its role in memory B cell function is unknown. IL-7R is likely to be functionally required for memory B cell self-renewal. However, it is unlikely to function at the level of stem cells, because the IL-7R protein is not expressed on Lt-HSC cell surfaces. A more straightforward explanation for its Lt-HSC expression is that IL-7R gene transcription lies downstream of a common self-renewal pathway. Alternatively, there are several shared transcripts that are

more likely to play a functional role in self-renewal. In particular, the signaling molecules mitogen-activated protein kinase 12 and PKC- $\zeta$  and the transcription factor Pou6f1 represent potentially convergent nodes in the network of self-renewal pathways.

Identification of these transcripts lends significant impetus for further testing of their functional relevance to hematopoietic and memory cell self-renewal. In particular, our data suggest that the polycomb complex that includes Bmi-1 is likely to function in memory B cell self-renewal in addition to its already reported role in hematopoietic stem cells. Given the role of polycomb genes in the maintenance of cellular memory of chromatin modification and transcriptional repression, these molecules are particularly intriguing candidates for functioning in immunologic memory. Further, it is worth considering the possibility that the separate pathways identified in our analysis might functionally converge within the cell. For instance, Bmi-1 itself has recently been shown to associate with and be phosphorylated by 3pK (mitogen-activated protein kinase AP kinase 3), which lies downstream of several mitogen-activated protein kinase pathways (44).

There has been a great deal of debate concerning the validity and reproducibility of defining a general molecular signature of stem cells (14, 27, 45, 46). Although “stemness” certainly requires an aspect of self-renewal, there are many additional functions and/or states that might be shared by the broad range of stem cells examined in the previous studies. Furthermore, it is not difficult to imagine arriving at similar phenotypes via divergent pathways, particularly within different lineages. Because memory T and memory B cells are descended from long-term and short-term HSC, we suggest that the focused comparisons presented herein provide unique insights into self-renewal within the hematopoietic system. Indeed, those genes shared between both memory populations were coordinately regulated in all three of the published HSC Affymetrix data sets we analyzed in Fig. 6. However, when the T cell data were considered in conjunction with previously published ES cell and neural stem cell data sets (14, 27), there was not a consistent enrichment of the shared transcripts in either of these two. This finding suggests that the molecular signature we defined may be restricted to the self-renewing cells of the hematopoietic system, a finding consistent with the published work of others showing conservation within, but not across, lineages (45, 46).

Our results have important implications beyond the identification of a self-renewal signature. For example, these shared transcripts are excellent candidates for those reactivated in the self-renewal program of leukemic stem cells (47–51). Indeed, an increase in the expression of the polycomb complex component Bmi-1 has been implicated in leukemogenesis (31, 52). Second, given the recent reports that memory CD8<sup>+</sup> T cell self-renewal is preferentially localized to the bone marrow (53), it is an intriguing possibility that memory T cells and hematopoietic stem cells may have partially overlapping niches within the marrow that support their self-renewal. Finally, the data provides a glimpse of the shared biochemical mechanisms with which hematopoietic cells undergo self-renewal.

## Methods

**T Cell Purification, RNA Processing, and Amplification.** T cells were sorted, and RNA was purified, amplified, and hybridized as described in ref. 15. Details for purification are given in *Supporting Methods*. Replicates included naïve (four), effector (three), and memory (five) populations.

**B Lineage Cell Purification, RNA Processing, and Amplification.** The purification strategy, ELISPOT assays, RNA processing method, and hybridization strategy are all detailed in *Supporting Methods*. Replicates included naïve (three), germinal center (three), plasma (four), and memory (four) populations.

**Hematopoietic Stem Cell Purification for Q-PCR Confirmation.** Lt-HSC, St-HSC, and LCP purification was performed by following the protocol described by Yang *et al.* (20). Details of purification strategy and Q-PCR methods are described in *Supporting Methods*.

**Statistical Methods.** Affymetrix image files (.cel) of the MGU74vA2-A chips from the stem cell and T cell data sets were collectively analyzed by using the AFFYLMGUI package developed by the open-source collaborative [www.bioconductor.org](http://www.bioconductor.org) (54). Data were background corrected, probe-level normalized and summarized by using the rma method (55). The rma method uses an improved algorithm for probe-level background correction, normalization, and summary that dramatically reduces observed statistical noise both in published control data sets (55–58) and among replicates within our own data (C.J.L., A.W.G., C.B., and D.M., unpublished results). Differential expression and false discovery rate-corrected *P* values were determined by using the LIMMA method (59). Affymetrix image files (.cel) of the 430.V2 chips from

the from the B cell data sets were collectively analyzed by using the AFFYLMGUI package as described above. These data sets were then linked at the probe level with the B and T cell data sets by using the published best match correlation files from Affymetrix.

We thank Amy Wagers for her helpful advice and suggestions; Natalia Ivanova, Ihor Lemischka, Miguel Ramalho-Santos, Douglas A. Melton, and Liheng Li for their help with data acquisition; Vanessa Tran, Quynh-Mai Pham, Vera Bruklich, and Tatyana Lipatova for assistance with mice; and Joyce LaVecchio and Grigoriy Losev for cell sorting. This work was supported by Joslin's Diabetes and Endocrinology Research Center Cores Grant 2 P30 DK36836-17, National Institutes of Health (NIH) Grant AI51530-01, and the W. T. Young Chair for Diabetes Research (to C.B. and D.M.) and NIH Grants R01HL058770, R01AI047457, and P01DK053074 (to I.L.W.). A.W.G. was supported by an Irvington Institute–Juvenile Diabetes Research Fellowship and Joslin's National Institutes of Health Training Grant 5 T32 DK007260-24. D.B. was funded by a Cancer Research Institute Fellowship. C.J.L. was funded by Brigham and Women's Pathology Department Training Grant T32 HL0762–18 and NIH Grant K08 AI063386.

1. Fearon, D. T., Manders, P. & Wagner, S. D. (2001) *Science* **293**, 248–250.
2. Kaech, S. M., Wherry, E. J. & Ahmed, R. (2002) *Nat. Rev. Immunol.* **2**, 251–262.
3. Lanzavecchia, A. & Sallusto, F. (2002) *Nat. Rev. Immunol.* **2**, 982–987.
4. Kondo, M., Wagers, A. J., Manz, M. G., Prohaska, S. S., Scherer, D. C., Beilhack, G. F., Shizuru, J. A. & Weissman, I. L. (2002) *Annu. Rev. Immunol.* **20**, 759–806.
5. Schluns, K. S. & Lefrancois, L. (2003) *Nat. Rev. Immunol.* **3**, 269–279.
6. Woodland, R. T. & Schmidt, M. R. (2005) *Semin. Immunol.* **17**, 209–217.
7. Macallan, D. C., Asquith, B., Irvine, A. J., Wallace, D. L., Worth, A., Ghattas, H., Zhang, Y., Griffin, G. E., Tough, D. F. & Beverley, P. C. (2003) *Eur. J. Immunol.* **33**, 2316–2326.
8. Sprent, J. (2003) *Microbes Infect.* **5**, 227–231.
9. Forster, I., Vieira, P. & Rajewsky, K. (1989) *Int. Immunol.* **1**, 321–331.
10. Schitteck, B. & Rajewsky, K. (1990) *Nature* **346**, 749–751.
11. Smith, K. G., Weiss, U., Rajewsky, K., Nossal, G. J. & Tarlinton, D. M. (1994) *Immunity* **1**, 803–813.
12. Slifka, M. K., Antia, R., Whitmire, J. K. & Ahmed, R. (1998) *Immunity* **8**, 363–372.
13. McHeyzer-Williams, L. J., Cool, M. & McHeyzer-Williams, M. G. (2000) *J. Exp. Med.* **191**, 1149–1166.
14. Ivanova, N. B., Dimos, J. T., Schaniel, C., Hackney, J. A., Moore, K. A. & Lemischka, I. R. (2002) *Science* **298**, 601–604.
15. Goldrath, A. W., Luckey, C. J., Park, R., Benoist, C. & Mathis, D. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 16885–16890.
16. Homann, D., Teyton, L. & Oldstone, M. B. (2001) *Nat. Med.* **7**, 913–919.
17. Wherry, E. J. & Ahmed, R. (2004) *J. Virol.* **78**, 5535–5545.
18. Kaech, S. M., Hemby, S., Kersh, E. & Ahmed, R. (2002) *Cell* **111**, 837–851.
19. Adolfsson, J., Borge, O. J., Bryder, D., Theilgaard-Monch, K., Astrand-Grundstrom, I., Sitnicka, E., Sasaki, Y. & Jacobsen, S. E. (2001) *Immunity* **15**, 659–669.
20. Yang, L., Bryder, D., Adolfsson, J., Nygren, J., Mansson, R., Sigvardsson, M. & Jacobsen, S. E. (2005) *Blood* **105**, 2717–2723.
21. Adolfsson, J., Mansson, R., Buza-Vidas, N., Hultquist, A., Liuba, K., Jensen, C. T., Bryder, D., Yang, L., Borge, O. J., Thoren, L. A., *et al.* (2005) *Cell* **121**, 295–306.
22. Christensen, J. L. & Weissman, I. L. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 14541–14546.
23. Forsberg, E. C., Prohaska, S. S., Katzman, S., Heffner, G. C., Stuart, J. M. & Weissman, I. L. (2005) *PLoS Genet.* **1**, e28.
24. Tersikh, A. V., Miyamoto, T., Chang, C., Diatchenko, L. & Weissman, I. L. (2003) *Blood* **102**, 94–101.
25. Park, I. K., He, Y., Lin, F., Laerum, O. D., Tian, Q., Bumgarner, R., Klug, C. A., Li, K., Kuhr, C., Doyle, M. J., Xie, T., *et al.* (2002) *Blood* **99**, 488–498.
26. Park, P. J., Cao, Y. A., Lee, S. Y., Kim, J. W., Chang, M. S., Hart, R. & Choi, S. (2004) *J. Biotechnol.* **112**, 225–245.
27. Ramalho-Santos, M., Yoon, S., Matsuzaki, Y., Mulligan, R. C. & Melton, D. A. (2002) *Science* **298**, 597–600.
28. Akashi, K., He, X., Chen, J., Iwasaki, H., Niu, C., Steenhard, B., Zhang, J., Haug, J. & Li, L. (2003) *Blood* **101**, 383–389.
29. Zhong, J. F., Zhao, Y., Sutton, S., Su, A., Zhan, Y., Zhu, L., Yan, C., Gallaher, T., Johnston, P. B., Anderson, W. F. & Cooke, M. P. (2005) *Proc. Natl. Acad. Sci. USA* **102**, 2448–2453.
30. Park, I. K., Qian, D., Kiel, M., Becker, M. W., Pihalja, M., Weissman, I. L., Morrison, S. J. & Clarke, M. F. (2003) *Nature* **423**, 302–305.
31. Lessard, J. & Sauvageau, G. (2003) *Nature* **423**, 255–260.
32. Molofsky, A. V., Pardal, R., Iwashita, T., Park, I. K., Clarke, M. F. & Morrison, S. J. (2003) *Nature* **425**, 962–967.
33. Liu, N., Phillips, T., Zhang, M., Wang, Y., Opferman, J. T., Shah, R. & Ashton-Rickardt, P. G. (2004) *Nat. Immunol.* **5**, 919–926.
34. Liu, N., Wang, Y. & Ashton-Rickardt, P. G. (2004) *FEBS Lett.* **569**, 49–53.
35. Zhang, Y., Schlossman, S. F., Edwards, R. A., Ou, C.-N., Gu, J. & Wu, M. X. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 878–883.
36. Zapata, J. M. & Reed, J. C. (2002) *Sci. STKE* **2002** (133), PE27.
37. Baud, V. & Karin, M. (2001) *Trends Cell Biol.* **11**, 372–377.
38. Dorrell, C., Takenaka, K., Minden, M. D., Hawley, R. G. & Dick, J. E. (2004) *Mol. Cell. Biol.* **24**, 6993–7002.
39. Qi, X., Li, T.-G., Hao, J., Hu, J., Wang, J., Simmons, H., Miura, S., Mishina, Y. & Zhao, G.-Q. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 6027–6032.
40. Burdon, T., Smith, A. & Savatier, P. (2002) *Trends Cell Biol.* **12**, 432–438.
41. Reya, T. & Clevers, H. (2005) *Nature* **434**, 843–850.
42. Mao, Z., Bonni, A., Xia, F., Nadal-Vicens, M. & Greenberg, M. E. (1999) *Science* **286**, 785–790.
43. Gaudilliere, B., Shi, Y. & Bonni, A. (2002) *J. Biol. Chem.* **277**, 46442–46446.
44. Voncken, J. W., Niessen, H., Neufeld, B., Rennefahrt, U., Dahlmans, V., Kubben, N., Holzer, B., Ludwig, S. & Rapp, U. R. (2005) *J. Biol. Chem.* **280**, 5178–5187.
45. Evsikov, A. V. & Solter, D. (2003) *Science* **302**, 393; author reply 393.
46. Fortunel, N. O., Otu, H. H., Ng, H. H., Chen, J., Mu, X., Chevassut, T., Li, X., Joseph, M., Bailey, C., Hatzfeld, J. A., *et al.* (2003) *Science* **302**, 393; author reply 393.
47. Lapidot, T., Sirard, C., Vormoor, J., Murdoch, B., Hoang, T., Caceres-Cortes, J., Minden, M., Paterson, B., Caligiuri, M. A. & Dick, J. E. (1994) *Nature* **367**, 645–648.
48. Sirard, C., Lapidot, T., Vormoor, J., Cashman, J. D., Doedens, M., Murdoch, B., Jamal, N., Messner, H., Addey, L., Minden, M., *et al.* (1996) *Blood* **87**, 1539–1548.
49. Jamieson, C. H., Ailles, L. E., Dylla, S. J., Muijtjens, M., Jones, C., Zehnder, J. L., Gotlib, J., Li, K., Manz, M. G., Keating, A., *et al.* (2004) *N. Engl. J. Med.* **351**, 657–667.
50. Passegue, E., Jamieson, C. H. M., Ailles, L. E. & Weissman, I. L. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 11842–11849.
51. Warner, J. K., Wang, J. C., Hope, K. J., Jin, L. & Dick, J. E. (2004) *Oncogene* **23**, 7164–7177.
52. Raaphorst, F. M. (2003) *Trends Immunol.* **24**, 522–524.
53. Di Rosa, F. & Pabst, R. (2005) *Trends Immunol.* **26**, 360–366.
54. Wettenhall, J. M. & Smyth, G. K. (2004) *Bioinformatics* **20**, 3705–3706.
55. Irizarry, R. A., Hobbs, B., Collin, F., Beazer-Barclay, Y. D., Antonellis, K. J., Scherf, U. & Speed, T. P. (2003) *Biostatistics* **4**, 249–264.
56. Irizarry, R. A., Bolstad, B. M., Collin, F., Cope, L. M., Hobbs, B. & Speed, T. P. (2003) *Nucleic Acids Res.* **31**, e15.
57. Bolstad, B. M., Irizarry, R. A., Astrand, M. & Speed, T. P. (2003) *Bioinformatics* **19**, 185–193.
58. Wu, Z. & Irizarry, R. A. (2004) *Nat. Biotechnol.* **22**, 656–658; author reply 658.
59. Smyth, G. K. (2004) *Statistical Applications in Genetics and Molecular Biology*, Vol. 3, Issue 1, article 3. Available at [www.bepress.com/sagmb/vol3/iss1/art3](http://www.bepress.com/sagmb/vol3/iss1/art3). Accessed January 23, 2006.

**Table 1.** Transcripts whose expression is coordinately regulated in memory T cells and Lt-HSC

				Positive fold change							
<b>Symbol</b>	<b>Name</b>	<b>Unigene</b>	<b>Affy ID</b>	<b>Lt v LCP</b>	<b>Lt v St</b>	<b>St v LCP</b>	<b>Tm v Tn</b>	<b>Tm v Te</b>	<b>Bm v Bn</b>	<b>Bm v Bgc</b>	<b>Bm v Bpl</b>
Mcoln2	Mucolin 2	Mm.116862	103686_at	5.43	1.83	2.97	2.14	2.04	0.80	1.64	0.51
Map3k8	Mitogen activated protein kinase kinase kinase 8	Mm.3275	1419208_at	5.35	2.13	2.51	2.17	1.49	0.76	5.75	9.39
Arl7	ADP-ribosylation factor-like 7	Mm.3388	1436512_at	4.79	1.65	2.91	1.37	2.20	0.71	0.97	0.90
NA	NA	Mm.29940	1448021_at	3.34	2.00	1.67	1.98	2.60	6.84	0.40	0.01
Ier3	Immediate early response 3	Mm.25613	161281_f_at	3.07	1.48	2.08	2.01	1.39	1.05	0.80	0.88
Trif-P	Ring finger protein 138	Mm.42154	1419368_a_at	3.05	1.64	1.86	2.06	1.40	1.11	1.24	1.99
Mel-18	Ring finger protein 110	Mm.2418	96961_at	2.62	1.91	1.37	1.39	1.56	1.04	0.83	1.04
Plekha5	Pleckstrin homology domain containing, family A member 5	Mm.80531	1425543_s_at	2.60	1.88	1.38	1.49	1.37	1.91	1.87	1.75
5Rik	RIKEN cDNA B930046C15 gene	Mm.204	1452731_x_at	2.30	1.65	1.39	1.59	2.19	3.80	3.76	5.22
Traf1	Tnf receptor-associated factor 1	Mm.12898	1423602_at	2.28	1.53	1.49	2.07	2.43	3.22	0.79	8.84
Rras	Harvey rat sarcoma oncogene, subgroup R	Mm.257	1418448_at	2.28	1.41	1.62	1.50	1.38	1.31	1.13	1.35
Prkce	RIKEN cDNA 5830406C15 gene	Mm.2013	1452878_at	2.25	1.69	1.33	1.73	1.44	0.74	2.85	6.13
Antxr2	Anthrax toxin receptor 2	Mm.24842	1426708_at	2.16	1.48	1.45	2.14	1.69	0.37	0.90	0.75
Nr1d2	Nuclear receptor subfamily 1, group D, member 2	Mm.26584	99076_at	2.13	1.52	1.41	2.28	1.68	0.85	5.78	2.81
Mapk12	Mitogen-activated protein kinase 12	Mm.38343	1449283_a_at	2.07	1.55	1.34	2.45	2.27	4.11	30.02	56.86
Lpin1	Lipin 1	Mm.28548	1418288_at	2.04	1.52	1.35	1.87	1.66	0.81	2.71	1.65
BC042396	cDNA sequence BC042396	Mm.24944	1455694_at	2.03	1.39	1.45	1.94	1.37	3.88	2.23	5.91
Rnf138	Ring finger protein 138	Mm.42154	1454064_a_at	2.01	1.81	1.11	2.55	2.04	0.71	1.11	1.89
U2af1-rs1	U2Af related sequence 1	Mm.14286	1449354_at	1.92	1.60	1.20	1.41	1.68	1.32	11.45	4.03
Txnip	Thioredoxin interacting protein	Mm.77432	1415996_at	1.92	1.96	0.98	1.39	2.77	0.95	8.83	3.02
Il7r	Interleukin 7 receptor	Mm.389	1448575_at	1.73	1.87	0.93	2.39	5.17	1.41	1.81	1.53
Tnfrsf1b	Tumor necrosis factor receptor superfamily, member 1b	Mm.2666	1418099_at	1.68	1.95	0.86	2.41	1.39	1.30	2.57	2.00
Dock9	Dedicator of cytokinesis 9	Mm.24477	1450932_s_at	1.54	1.57	0.98	1.92	2.11	2.22	1.34	10.21

				Negative fold change							
<b>Symbol</b>	<b>Name</b>	<b>Unigene</b>	<b>Affy ID</b>	<b>Lt v LCP</b>	<b>Lt v St</b>	<b>St v LCP</b>	<b>Tm v Tn</b>	<b>Tm v Te</b>	<b>Bm v Bn</b>	<b>Bn v Bgc</b>	<b>Bm v Bpl</b>
Plac8	Placenta-specific 8	Mm.34609	1451335_at	5.17	3.43	1.51	1.80	1.85	0.76	0.01	0.17
Ezh2	Enhancer of zeste homolog 2 (Drosophila)	Mm.4303	1416544_at	3.73	2.00	1.86	1.79	6.36	1.05	9.22	3.59
Trim59	Tripartite motif-containing 59	Mm.41535	1416118_at	3.53	1.80	1.96	1.78	2.08	2.46	2.42	0.82
Kif11	Kinesin family member 11	Mm.42203	1435306_a_at	3.51	3.89	0.90	1.37	7.16	1.13	12.89	8.27
Anp32e	E	Mm.218657	1451356_at	2.87	1.49	1.93	1.38	2.31	1.53	3.10	0.91
Coq7	Demethyl-Q 7	Mm.20634	1416665_at	2.62	1.55	1.69	1.63	1.74	1.60	3.42	4.98
Ywhah	Eta polypeptide	Mm.3308	1416004_at	2.57	1.52	1.68	1.39	2.85	1.51	2.78	1.70
Rasl2-9	RAS-like, family 2, locus 9	Mm.103632	1422656_at	2.41	2.14	1.12	1.38	1.74	1.14	2.13	1.09
Lbr	Lamin B receptor	Mm.4538	1415829_at	2.38	1.41	1.68	1.39	1.72	1.35	1.27	2.08
Smc4l1	SMC4 structural maintenance of chromosomes 4-like 1 (yeast)	Mm.206841	1452197_at	2.28	1.91	1.19	3.66	2.57	0.93	2.01	1.35
Eif1ay	Eukaryotic translation initiation factor 1A, Y-linked	Mm.65264	1419736_a_at	2.23	1.40	1.60	1.41	2.06	1.81	3.68	3.71
Kif20a	Kinesin family member 20A	Mm.196638	1449207_a_at	2.16	2.99	0.72	1.40	6.63	1.12	51.25	18.96
Cbx3	Chromobox homolog 3 (Drosophila HP1 gamma)	Mm.28148	1448504_a_at	1.83	1.53	1.19	1.43	1.67	1.38	1.81	1.78
Ranbp1	RAN binding protein 1	Mm.3752	98573_f_at	1.82	1.46	1.25	1.74	1.47	2.69	4.72	3.75
Tubb5	Tubulin, beta 5	Mm.1703	1416256_a_at	1.79	2.58	0.69	1.98	1.91	1.32	2.22	2.93
Wtap	Wilms' tumour 1-associating protein	Mm.25154	1454805_at	1.76	1.50	1.17	1.41	1.56	1.76	1.50	1.01
Hdgf	Hepatoma-derived growth factor	Mm.195277	1419964_s_at	1.75	1.62	1.08	1.59	1.96	1.35	1.67	2.66
Tuba2	Tubulin, alpha 2	Mm.196396	1423846_x_at	1.73	2.22	0.78	1.49	1.98	1.09	1.72	1.25
Tuba6	Tubulin, alpha 6	Mm.88212	101543_f_at	1.52	1.91	0.80	1.99	1.47	0.78	0.72	0.73
Tuba1	Tubulin, alpha 1	Mm.196396	1418884_x_at	1.49	2.06	0.72	1.51	2.04	1.18	1.79	0.71
Hnrpa2b1	Heterogeneous nuclear ribonucleoprotein A2/B1	Mm.16767	1420365_a_at	1.43	1.58	0.91	1.67	1.95	1.61	1.94	1.64

Lt v LCP is the fold change between Lt-HSC and LCP. Lt v St is the fold change between Lt-HSC and St-HSC. Tm v Tn is the fold change between memory and naive CD8+ T cells. Tm v Te is the fold change between memory and effector CD8+ T cells. Bm v Bn is the fold change between memory and naive B cells. Bm v Bgc is the fold change between memory and germinal center B cells. Bm v Bpl is the fold change between memory and plasma B cells.

**Table 2.** Transcripts whose expression is coordinately up-regulated in memory B cells and Lt-HSC


**Up-regulated transcripts**

				Positive fold change							
Symbol	Name	Unigene	Affy ID	Lt v LCP	Lt v St	St v LCP	Tm v Tn	Tm v Te	Bm v Bn	Bn v Bgc	Bm v Bpl
5730557B15Rik	RIKEN cDNA 5730557B15 gene	Mm.27619	1453287_at	10.70	3.14	3.41	1.13	1.18	8.29	5.01	52.58
Gem	GTP binding protein (gene overexpressed in skeletal muscle	Mm.4362	1426063_a_at	7.41	2.38	3.12	1.06	0.34	1.38	1.58	2.01
C79248	Expressed sequence C79248	Mm.153895	1442745_x_at	4.96	1.64	3.03	1.03	1.01	1.64	1.79	4.05
Ccl3	Chemokine (C-C motif) ligand 3	Mm.1282	1419561_at	4.08	1.89	2.16	1.64	0.22	1.99	1.75	1.63
Bteb1	Basic transcription element binding protein 1	Mm.183017	1428289_at	4.00	2.31	1.73	1.13	0.85	1.58	2.14	4.48
AI839402	Expressed sequence AI839402	Mm.202684	1435885_s_at	3.94	2.41	1.64	1.20	0.87	1.65	2.07	2.68
AI646383	Expressed sequence AI646383	Mm.2304	1459922_at	3.48	1.99	1.75	1.00	1.08	3.15	2.21	4.26
Zfp612	Zinc finger protein 612	Mm.87487	1427104_at	3.43	3.01	1.14	1.16	1.18	7.24	4.71	7.68
Lilrb4	Leukocyte immunoglobulin-like receptor, subfamily B, memb	Mm.34408	1420394_s_at	3.27	2.38	1.38	4.35	0.47	33.83	51.16	36.84
NA	NA	Mm.182434	1424609_a_at	3.23	2.25	1.43	0.72	1.78	3.16	4.98	4.88
li	la-associated invariant chain	Mm.7043	1425519_a_at	3.20	1.96	1.63	2.00	0.67	1.79	2.87	2.73
Itpr4	Inositol 1,4,5-triphosphate receptor 4	Mm.7800	1427287_s_at	3.10	1.53	2.03	0.70	2.00	2.17	2.28	9.73
Gabbr1	Gamma-aminobutyric acid (GABA-B) receptor, 1	Mm.32191	1455021_at	3.07	2.03	1.53	0.89	1.31	2.92	3.00	4.00
Tcf4	Transcription factor 4	Mm.4269	1416723_at	3.07	1.55	1.99	0.81	0.92	1.94	3.45	1.74
D8Ert4325e	DNA segment, Chr 8, ERATO Doi 325, expressed	Mm.2875	1437205_at	2.97	1.55	1.91	1.11	1.87	1.93	3.75	2.35
Ptpn1	Protein tyrosine phosphatase, non-receptor type 1	Mm.2668	1417068_a_at	2.89	1.50	1.92	0.98	1.25	1.59	1.57	4.60
Mef2a	Myocyte enhancer factor 2A	Mm.87279	1427186_a_at	2.87	1.77	1.62	0.69	0.81	2.49	6.22	3.81
Ube1l	Ubiquitin-activating enzyme E1-like	Mm.1183	1426970_a_at	2.83	1.81	1.57	0.73	1.56	1.42	3.03	5.79
Gimap4	GTPase, IMAP family member 4	Mm.28395	1424375_s_at	2.77	3.05	0.91	1.09	0.95	1.83	10.42	7.28
C230027N18Rik	RIKEN cDNA C230027N18 gene	Mm.32738	1437111_at	2.77	1.91	1.45	0.88	0.95	4.87	7.28	6.57
Ifi205	Interferon activated gene 205	Mm.215120	1452348_s_at	2.73	1.68	1.63	0.63	1.13	2.80	45.80	7.51
Ahr	Aryl-hydrocarbon receptor	Mm.4452	1422631_at	2.71	2.20	1.23	1.14	1.25	1.97	4.54	1.69
Pbx3	Pre B-cell leukemia transcription factor 3	Mm.137604	1447640_s_at	2.69	2.03	1.33	1.80	0.65	4.24	2.50	6.39
Smarca2	SWI/SNF related, matrix associated, actin dependent regulat	Mm.12184	1430526_a_at	2.64	3.48	0.76	0.36	1.04	1.61	6.33	4.61
Plekha5	Pleckstrin homology domain containing, family A member 5	Mm.80531	1425543_s_at	2.60	1.88	1.38	1.49	1.37	1.91	1.87	1.75
5730454B08Rik	RIKEN cDNA 5730454B08 gene	Mm.193073	1426361_at	2.57	1.44	1.78	0.92	1.91	1.63	4.11	2.23
Wdr26	WD repeat domain 26	Mm.21126	1438234_at	2.53	1.66	1.52	0.68	1.13	2.20	2.77	3.62
Akap8l	A kinase (PRKA) anchor protein 8-like	Mm.9590	1417734_at	2.46	1.42	1.74	0.92	2.16	1.61	3.74	2.43
Impact	Imprinted and ancient	Mm.8154	1415911_at	2.45	1.76	1.39	1.08	1.14	1.92	2.34	1.70
Idb2	Inhibitor of DNA binding 2	Mm.1466	1435176_a_at	2.45	2.95	0.83	9.45	0.54	8.44	1.85	4.44
Dusp1	Dual specificity phosphatase 1	Mm.2404	1448830_at	2.31	1.55	1.49	1.60	0.45	1.84	1.71	2.83
B930046C15Rik	RIKEN cDNA B930046C15 gene	Mm.204	1452731_x_at	2.30	1.65	1.39	1.59	2.19	3.80	3.76	5.22
Capg	Capping protein (actin filament), gelsolin-like	Mm.18626	1450355_a_at	2.27	1.49	1.52	2.23	0.60	1.48	33.12	2.64
Csad	Cysteine sulfonic acid decarboxylase	Mm.41853	1427981_a_at	2.19	1.44	1.52	0.88	1.12	1.50	2.35	1.42
Mapk12	Mitogen-activated protein kinase 12	Mm.38343	1449283_a_at	2.07	1.55	1.34	2.45	2.27	4.11	30.02	56.86
Mef2d	Myocyte enhancer factor 2D	Mm.28184	1434487_at	2.04	1.70	1.20	0.91	1.21	1.47	1.78	1.89
BC042396	CDNA sequence BC042396	Mm.24944	1455694_at	2.03	1.39	1.45	1.94	1.37	3.88	2.23	5.91
Abca1	ATP-binding cassette, sub-family A (ABC1), member 1	Mm.369	1421840_at	1.98	2.23	0.89	1.01	2.53	1.45	13.78	9.65
Phr1	Pam, highwire, rpm 1	Mm.6478	1434937_at	1.93	2.07	0.93	0.62	1.50	3.15	2.93	7.36
Phxr4	Per-hexamer repeat gene 4	Mm.41972	1422272_at	1.93	1.48	1.30	0.80	1.64	10.00	16.20	10.94
Klf7	Kruppel-like factor 7 (ubiquitous)	Mm.29466	1419354_at	1.91	1.62	1.18	0.70	1.44	1.69	2.97	3.90
Gdap10	Ganglioside-induced differentiation-associated-protein 10	Mm.12950	1420342_at	1.90	1.96	0.97	0.72	0.85	2.06	6.53	9.46
Klf7	Kruppel-like factor 7 (ubiquitous)	Mm.44063	1437917_at	1.76	1.77	0.99	0.70	1.68	1.42	3.99	6.87
Tcf12	Transcription factor 12	Mm.45532	1430195_at	1.76	1.72	1.02	0.70	1.44	2.52	1.72	6.67
Prkwnk1	Protein kinase, lysine deficient 1	Mm.27341	1436746_at	1.75	1.47	1.20	0.84	1.02	1.67	6.45	2.27
Atp7a	ATPase, Cu++ transporting, alpha polypeptide	Mm.14926	1418774_a_at	1.73	1.39	1.24	1.03	1.23	1.72	2.73	1.99
Il7r	Interleukin 7 receptor	Mm.389	1448575_at	1.73	1.87	0.93	2.39	5.17	1.41	1.81	1.53
D2Ert463e	DNA segment, Chr 2, ERATO Doi 63, expressed	Mm.24965	1420253_at	1.73	1.53	1.13	0.96	0.85	1.83	1.95	1.96
Phxr1	Per-hexamer repeat gene 1	Mm.41987	1421800_at	1.72	1.57	1.10	1.03	1.07	1.69	2.10	4.76
B930046C15Rik	RIKEN cDNA B930046C15 gene	Mm.204	1452731_x_at	1.69	1.37	1.23	1.27	1.52	3.80	3.76	5.22
Chd7	Chromodomain helicase DNA binding protein 7	Mm.21099	1448026_at	1.63	2.22	0.74	0.82	0.93	2.41	4.33	3.86
Lmo2	LIM domain only 2	Mm.29266	1454086_a_at	1.61	1.45	1.11	1.15	0.76	1.47	3.20	96.34
Nedd4	Neural precursor cell expressed, developmentally down-regu	Mm.16553	1450431_a_at	1.58	1.53	1.04	0.89	1.61	3.00	2.06	1.39
Il6ra	Interleukin 6 receptor, alpha	Mm.2856	1452416_at	1.56	1.42	1.10	0.55	2.41	2.91	6.12	2.59
D7Ert4183e	DNA segment, Chr 7, ERATO Doi 183, expressed	Mm.24993	1449755_at	1.54	1.55	0.99	0.99	1.37	2.64	1.48	3.59
Hnrpa2b1	Heterogeneous nuclear ribonucleoprotein A2/B1	Mm.16767	1433829_a_at	1.50	1.83	0.82	0.77	0.74	1.90	1.55	1.39
LOC216024	Similar to heterogeneous nuclear ribonucleoprotein H3, isofo	Mm.28070	1455491_at	1.45	1.43	1.02	1.01	1.18	1.83	1.52	2.29
AI451896	Expressed sequence AI451896	Mm.25583	1459917_at	1.41	1.78	0.79	1.32	2.14	1.52	3.55	7.51
AA517132	Expressed sequence AA517132	Mm.203866	1452217_at	1.41	1.66	0.85	6.23	0.86	8.12	36.06	20.89
Prkwnk1	Protein kinase, lysine deficient 1	Mm.21773	1433676_at	1.38	1.74	0.79	0.82	1.09	1.57	1.51	2.14
NA	NA	Mm.1065	1459920_at	1.38	1.71	0.80	1.25	2.06	1.75	1.88	2.93
Bmi1	B lymphoma Mo-MLV insertion region 1	Mm.7719	1448733_at	1.05	0.55	0.58	1.59	1.08	1.67	12.99	2.44


See Table 1 legend for explanation of abbreviations.

**Table 3. Transcripts whose expression is coordinately down-regulated in memory B cells and Lt-HSC**

Down-regulated transcripts		Negative fold change									
Symbol	Name	Unigene	AffID	Lt v LCP	Lt v St	St v LCP	Tm v Tn	Tm v Te	Bm v Bn	Bn v Boc	Bm v Bnl
Ap3s1	Adaptor-related protein complex 3, sigma 1 subunit	Mm.27171	1422593_at	10.93	1.86	5.90	0.90	1.50	2.20	4.91	8.37
Fignl1	Fidgletin-like 1	Mm.20315	1422430_at	10.41	2.64	3.94	1.02	5.10	2.05	28.31	21.30
Kpna2	Karyopherin (importin) alpha 2	Mm.12508	1415860_at	8.34	2.68	3.12	1.36	5.94	1.47	4.76	4.74
Usp1	Ubiquitin specific protease 1	Mm.27496	1451080_at	6.59	2.45	2.68	0.97	2.25	1.54	1.99	2.01
Plk4	Polo-like kinase 4 (Drosophila)	Mm.198533	1419838_s_at	5.21	2.77	1.88	1.07	3.20	1.42	7.92	5.45
Tacc3	Transforming, acidic coiled-coil containing protein 3	Mm.27836	1417450_a_at	4.76	3.97	1.19	0.90	5.62	1.75	19.41	6.76
Psm1	Proteasome (prosome, macropain) subunit, alpha type 1	Mm.30097	1415695_at	4.50	1.40	3.23	1.25	2.46	1.90	2.62	4.07
Psm1	Proteasome (prosome, macropain) subunit, alpha type 1	Mm.30097	1415695_at	4.50	1.40	3.23	1.25	2.46	1.90	2.62	4.07
Rfc4	Replication factor C (activator 1) 4	Mm.18876	1424321_at	4.41	2.53	1.74	0.91	2.30	2.77	14.11	12.87
Tmem14c	Transmembrane protein 14C	Mm.30005	1416479_a_at	4.29	1.61	2.66	1.16	1.65	1.51	2.73	3.18
Chek1	Checkpoint kinase 1 homolog (S. pombe)	Mm.197875	1449708_s_at	4.29	2.53	1.70	0.96	2.42	1.48	59.10	29.23
Cks1b	CDC28 protein kinase 1b	Mm.3049	1416698_a_at	4.23	2.99	1.42	0.98	5.21	2.28	72.02	84.30
Mcm4	Minichromosome maintenance deficient 4 homolog (S. cerevisiae)	Mm.1500	1416214_at	3.94	2.39	1.64	1.01	2.08	1.73	4.14	3.65
Fsmt1	Phosphoserine aminotransferase 1	Mm.29902	1451064_a_at	3.92	2.50	1.57	1.33	2.93	1.72	10.43	13.82
Rolt	SoxL/ZfSox9 leucine zipper binding protein in testis	Mm.2934	1452892_at	3.89	2.79	1.40	0.84	1.62	1.41	87.55	44.16
Rbl1	Retinoblastoma-like 1 (p107)	Mm.2994	1424156_at	3.84	1.60	2.41	1.02	1.45	1.39	2.56	1.70
Mrpl18	Mitochondrial ribosomal protein L18	Mm.21356	1448373_at	3.76	2.27	1.66	1.05	1.65	1.72	3.61	4.47
Mrpl36	Mitochondrial ribosomal protein L36	Mm.26775	1422819_at	3.73	1.75	2.13	1.09	1.18	1.52	1.48	2.72
181003N24Rik	RIKEN cDNA 181003N24 gene	Mm.27831	1450735_at	3.56	1.56	2.28	0.96	0.97	1.39	1.76	2.01
Mrpl18	Mitochondrial ribosomal protein L18	Mm.21356	1448373_at	3.43	2.01	1.71	1.03	1.44	1.72	3.61	4.47
Cdc20	Cell division cycle 20 homolog (S. cerevisiae)	Mm.29931	1416664_at	3.32	3.29	1.01	1.05	5.58	1.63	14.53	30.24
Rrm1	Ribonucleotide reductase M1	Mm.656	1415878_at	3.23	2.38	1.36	0.84	3.32	1.89	8.32	4.72
Topbp1	Topoisomerase (DNA) II beta binding protein	Mm.1687	1452241_at	3.16	1.92	1.64	1.12	1.77	1.57	3.15	2.85
Cops5	COP9 (constitutive photomorphogenic) homolog, subunit 5 (Mm.2472)	Mm.2472	1460171_at	3.07	1.48	2.08	1.24	1.65	1.39	1.76	4.45
Tipin	Timeless interacting protein	Mm.196219	1426612_at	2.99	2.04	1.46	0.91	2.57	1.56	8.79	6.28
Prim1	DNA primase, p49 subunit	Mm.2903	1449061_a_at	2.97	2.46	1.20	1.00	2.68	1.63	12.13	3.72
Mcm5	Minichromosome maintenance deficient 5, cell division cycle	Mm.5048	1415945_at	2.97	2.81	1.06	0.89	2.83	2.42	11.27	14.87
Nup62	Nucleoporin 62	Mm.22687	1415926_at	2.91	1.93	1.51	0.97	1.61	1.54	1.93	1.87
1810045K17Rik	RIKEN cDNA 1810045K17 gene	Mm.28917	1418899_at	2.71	1.66	1.64	0.82	1.25	1.38	2.41	8.56
Mrps14	Mitochondrial ribosomal protein S14	Mm.29599	1420489_at	2.71	1.39	1.94	1.00	1.06	1.37	2.78	5.12
Coq7	Demethyl-Q 7	Mm.20634	1416665_at	2.62	1.55	1.69	1.63	1.74	1.60	3.42	4.98
Ywhah	Tyrosine 3-monooxygenase/tryptophan 5-monooxygenase ac	Mm.3308	1416004_at	2.57	1.52	1.68	1.39	2.85	1.51	2.57	1.70
Mcm2	Minichromosome maintenance deficient 2 mitotin (S. cerevisiae)	Mm.16711	1448777_at	2.57	2.00	1.29	0.99	1.63	1.97	11.13	12.61
Mcm7	Minichromosome maintenance deficient 7 (S. cerevisiae)	Mm.18923	1416030_a_at	2.55	3.07	0.83	0.99	1.91	2.88	9.04	13.04
Nduaf2	NADH dehydrogenase (ubiquinone) flavoprotein 2	Mm.2964	1452892_at	2.49	1.70	1.36	1.11	1.26	1.44	1.85	2.48
Bra1	Breast cancer 1	Mm.1889	1424629_at	2.48	2.48	1.00	1.07	2.38	1.49	13.92	12.09
Asf1b	ASF1 anti-silencing function 1 homolog B (S. cerevisiae)	Mm.29680	1423714_at	2.46	1.92	1.28	1.08	3.68	1.37	17.19	10.42
Bub3	Budding uninhibited by benzimidazoles 3 homolog (S. cerevisiae)	Mm.927	1416815_s_at	2.46	1.77	1.39	1.25	1.76	1.43	2.30	1.87
Hmgb1	High mobility group box 1	Mm.16421	1416176_at	2.45	1.72	1.42	1.06	1.74	1.47	2.28	3.07
Ctps	Cytidine 5'-triphosphate synthase	Mm.1815	1416563_at	2.45	1.49	1.64	1.53	1.23	1.74	3.80	5.21
Psm5a	Proteasome (prosome, macropain) subunit, alpha type 5	Mm.2287	1424681_a_at	2.35	1.55	1.54	1.62	1.54	1.62	2.22	2.48
Hat1	Histone aminotransferase 1	Mm.28421	1428061_at	2.35	1.62	1.44	1.07	1.56	1.98	6.36	3.08
AW547477	Expressed sequence AW547477	Mm.124	1426349_s_at	2.33	2.04	1.14	0.99	2.13	1.68	2.36	2.69
Mrpl16	Mitochondrial ribosomal protein L16	Mm.41390	1450880_at	2.31	1.50	1.54	0.85	1.07	1.89	1.47	2.47
Mcm3	Minichromosome maintenance deficient 3 (S. cerevisiae)	Mm.200664	1420028_s_at	2.30	2.50	0.92	1.07	2.73	1.92	9.77	6.58
Hat1	Histone aminotransferase 1	Mm.28421	1428061_at	2.27	1.43	1.59	0.98	1.55	1.98	6.36	3.08
Eif1ay	Eukaryotic translation initiation factor 1A, Y-linked	Mm.65264	1419736_a_at	2.23	1.40	1.60	1.41	2.06	1.81	3.68	3.71
Phf5a	PHD finger protein 5A	Mm.29503	1424170_at	2.22	1.58	1.41	1.25	1.28	1.95	3.42	4.29
Rad51	RAD51 homolog (S. cerevisiae)	Mm.231	1418281_at	2.22	1.95	1.14	0.92	2.60	1.38	35.10	25.14
Mthfd2	Methylenetetrahydrofolate dehydrogenase (NAD+ dependent)	Mm.443	1419253_at	2.20	2.45	0.90	0.85	1.74	1.61	6.97	4.71
2310042G06Rik	RIKEN cDNA 2310042G06 gene	Mm.182294	1448543_at	2.20	1.38	1.00	1.74	1.84	3.08	3.29	3.29
2700085E05Rik	RIKEN cDNA 2700085E05 gene	Mm.38521	1428421_a_at	2.19	1.37	1.60	1.00	1.19	1.46	2.01	1.51
Dck	Deoxycytidine kinase	Mm.3446	1449176_a_at	2.19	1.45	1.51	1.19	2.00	1.80	4.58	2.00
Rfc5	Replication factor C (activator 1) 5	Mm.27997	1452917_at	2.17	2.23	0.97	0.95	2.07	1.86	4.39	5.53
AA959742	Expressed sequence AA959742	Mm.28992	1416294_at	2.17	1.50	1.70	0.92	1.69	1.69	2.48	18.51
Epha5	Eph receptor A5	Mm.91	1433507_at	2.14	1.73	1.40	1.11	5.56	1.44	2.48	2.88
Cpsf5	Cleavage and polyadenylation specific factor 5	Mm.28961	1417681_at	2.14	1.60	1.34	1.19	1.72	1.37	2.49	1.47
Atp5o	ATP synthase, H+ transporting, mitochondrial F1 complex, O	Mm.41	1416278_a_at	2.14	1.88	1.14	1.09	1.24	1.54	2.58	4.56
1810063B05Rik	RIKEN cDNA 1810063B05 gene	Mm.31946	1435864_a_at	2.13	1.43	1.48	0.97	0.91	1.45	1.83	1.57
Thoc4	THO complex 4	Mm.1886	1417724_at	2.11	1.58	1.34	1.17	2.20	1.88	1.68	2.67
Cdca4	Cell division cycle associated 4	Mm.28595	1423683_at	2.11	1.68	1.26	1.14	1.10	1.97	2.12	1.42
Dnmt1	DNA methyltransferase (cytosine-5) 1	Mm.7814	1422946_a_at	2.10	1.88	1.12	0.92	1.40	1.47	3.62	2.31
Umps	Uridine monophosphate synthetase	Mm.202767	1434859_at	2.08	1.46	1.43	1.06	1.13	1.63	5.40	3.53
Ndufs6	NADH dehydrogenase (ubiquinone) Fe-S protein 6	Mm.29897	1433603_at	2.06	1.49	1.38	0.83	2.09	1.49	2.58	4.34
Rangap1	RAN GTPase activating protein 1	Mm.3833	1423749_s_at	2.01	1.92	1.05	1.16	1.47	1.61	3.24	4.55
Ndufab1	NADH dehydrogenase (ubiquinone) 1, alpha/beta subcomplex	Mm.3014	1428159_s_at	1.99	1.83	1.09	1.06	1.04	1.46	2.40	2.77
Elavl1	ELAV (embryonic lethal, abnormal vision, Drosophila)-like 1 (Mm.119162)	Mm.119162	1448151_at	1.98	1.38	1.44	1.58	1.23	1.48	2.03	1.51
Ris2	Retroviral integration site 2	Mm.21873	1424143_a_at	1.97	1.83	1.00	0.89	1.12	2.14	1.80	1.72
1500040F11Rik	RIKEN cDNA 1500040F11 gene	Mm.196343	1430326_s_at	1.95	2.04	0.96	1.15	1.31	1.57	3.13	6.10
Timm23	Translocase of inner mitochondrial membrane 23 homolog (y	Mm.24565	1416485_at	1.93	1.38	1.40	1.25	1.27	1.50	1.96	2.60
Fxn	Frataxin	Mm.7319	1427282_a_at	1.93	1.71	1.13	1.05	1.30	1.56	4.27	3.90
Fen1	Flap structure specific endonuclease 1	Mm.2952	1421731_a_at	1.93	2.30	0.84	0.80	1.91	1.56	4.10	9.87
Shmt1	Serine hydroxymethyl transferase 1 (soluble)	Mm.3379	1425179_at	1.90	1.91	0.99	1.00	0.95	1.59	3.59	4.12
Psmb5	Proteasome (prosome, macropain) subunit, beta type 5	Mm.8911	1415676_a_at	1.88	1.74	1.08	1.33	1.30	1.63	2.17	5.31
Psmb5	Proteasome (prosome, macropain) subunit, beta type 5	Mm.8911	1415676_a_at	1.88	1.74	1.08	1.33	1.30	1.63	2.17	5.31
Mcm3	Minichromosome maintenance deficient 3 (S. cerevisiae)	Mm.200664	1420028_s_at	1.86	2.41	0.97	0.97	2.14	1.92	9.77	6.58
Rars	Arginyl-tRNA synthetase	Mm.27526	1416312_at	1.85	1.51	1.22	0.95	1.07	1.56	1.74	3.79
Cbx3	Chromobox homolog 3 (Drosophila HP1 gamma)	Mm.28148	1448504_a_at	1.83	1.53	1.19	1.43	1.67	1.38	1.81	1.78
Psmb3	Proteasome (prosome, macropain) subunit, beta type 3	Mm.21874	1460198_a_at	1.82	1.74	1.05	1.20	1.59	2.07	1.83	1.83
Prim2	DNA primase, p58 subunit	Mm.27705	1418036_at	1.82	2.57	0.71	0.86	1.46	1.82	5.01	4.51
Stmn1	Stathmin 1	Mm.28479	1415849_s_at	1.80	2.07	0.87	0.95	7.31	2.83	51.42	26.76
na	Gene model 1921, (NCBI)	Mm.1710	1426475_at	1.79	1.41	1.27	1.10	1.60	1.57	1.71	5.99
1700013H19Rik	RIKEN cDNA 1700013H19 gene	Mm.27777	1429270_a_at	1.79	2.13	0.84	0.92	1.82	3.06	14.89	1.72
Sf3a3	Splicing factor 3a, subunit 3	Mm.25779	1423811_at	1.79	1.58	1.13	1.20	1.02	2.17	2.48	1.87
Timeless	Timeless homolog (Drosophila)	Mm.6458	1417586_at	1.76	1.60	1.10	1.13	1.25	3.51	7.80	8.44
Cops3	COP9 (constitutive photomorphogenic) homolog, subunit 3 (Mm.40)	Mm.40	1416678_at	1.75	1.73</						



**Top 75 Memory Specific Genes**  
Memory vs. Naive  $\log_2(\text{Fold-change})$



**Corresponding Fold-change Kaech et al.**  
Memory vs. Naive  $\log_2(\text{Fold-change})$

